

AD-A137 511

ON THE THEORY OF ELECTRON H-ATOM AND PROTON TRANSFERS
(U) CALIFORNIA INST OF TECH PASADENA ARTHUR AMOS NOYES
LAB OF CHEMICAL PHYSICS R A MARCUS 17 JAN 84 TR-14
N00014-79-C-0009

1/1


UNCLASSIFIED

F/G 7/4

NL



END
- 1
FILMED
3
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

12

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 14	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ON THE THEORY OF ELECTRON, H-ATOM AND PROTON TRANSFERS		5. TYPE OF REPORT & PERIOD COVERED Technical Report 14
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Rudolph A. Marcus		8. CONTRACT OR GRANT NUMBER(s) N00014-79-C-0009
9. PERFORMING ORGANIZATION NAME AND ADDRESS Noyes Laboratory of Chemical Physics California Institute of Technology Pasadena, California 91125		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Task No. NR 359-702
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE January 17, 1984
		13. NUMBER OF PAGES 20
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) <div style="text-align: right;">DTIC ELECTRIC FEB 06 1984 E</div>		
18. SUPPLEMENTARY NOTES Prepared for publication in "Proceedings Electrochemical Society Symposium on the Chemistry and Physics of Electrocatalysis", J. D. E. McIntyre and M. J. Weaver, eds.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Electron transfers; proton transfers; atom transfers.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The initial part of this article discusses some recent experiments on elec- tron transfer reactions and comments on the related theory, including the frequency of solvent fluctuations. Also described are recent advances with Babamov and Lopez on the dynamics of H-atom transfer reactions. We then outline the application of the basic idea to H and H⁺ transfers in complex systems and in condensed phases (fast H-coordinates, slow remaining coordinates). Similarities and differences between electron transfers and		

AD A1 37511

DTIC FILE COPY

DD FORM 1473 1 JAN 73 EDITION OF 1 NOV 65 IS OBSOLETE

84 02 06 013

Unclassified
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

H or H⁺ transfers are next described. The relation of the present theory of such transfers to conventional transition state theory is discussed.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



OFFICE OF NAVAL RESEARCH

Contract N00014-79-C-0009

Task No. NR 359-702

TECHNICAL REPORT NO. 14

On the Theory of Electron, H-Atom and Proton Transfers

by

Rudolph A. Marcus

Prepared for Publication

in the

**Proceedings Electrochemical Society Symposium on the
Chemistry and Physics of Electrocatalysis**

**California Institute of Technology
Department of Chemistry
Pasadena, California 91125**

**Reproduction in whole or in part is permitted for
any purpose of the United States Government**

**This document has been approved for public release
and sale; its distribution is unlimited**

ON THE THEORY OF ELECTRON, H-ATOM
AND PROTON TRANSFERS* *

R. A. Marcus
Arthur Amos Noyes Laboratory of Chemical Physics
California Institute of Technology
Pasadena, California 91125

$H(+)$

$H(-)$

The initial part of this article discusses some recent experiments on electron transfer reactions and comments on the related theory, including the frequency of solvent fluctuations. Also described are recent advances with Babamov and Lopez on the dynamics of H-atom transfer reactions. We then outline the application of the basic idea to H and H^+ transfers in complex systems and in condensed phases (fast H-coordinates, slow remaining coordinates). Similarities and differences between electron transfers and H or H^+ transfers are next described. The relation of the present theory of such transfers to conventional transition state theory is discussed.

-1-

Presented at the Symposium on "The Chemistry and Physics of Electro-catalysis", Electrochemical Society meeting in San Francisco, May 8-13, 1983.

I. Outline

There have been a number of recent results for electron transfer reactions (described in Sec. II) and a recent development in the theory of light atom transfer (e. g., H, D) reactions by Babamov and Lopez in our group (described in Sec. III) for collinear systems such as $I + HI \rightarrow IH + I$. The latter theory is based on a "Born-Oppenheimer" separation in which the I's are the slow particles and H the fast ones. We then indicate in Sec. III how this concept is extended to more complicated processes, such as those occurring in condensed phases. We also note the breakdown of simple Franck-Condon calculations when applied to atom transfers. In Sec. IV some similarities and differences in the transfer of electrons between reactants, as compared with the transfer of light nuclei such as H and H^+ , are described. The relation of the theory of all three to conventional transition state theory is discussed.

II. Recent Developments in Electron Transfers

This part of the paper discusses some recent work in this field. The topics we consider are

1. electronic matrix elements in electron transfers; nonadiabaticity vs adiabaticity;
2. absolute rate calculations of electron transfer rate constants;
3. solvent effects, both when solvent relaxation is rate controlling and when it is not; an expression for the rate of formation of solvent fluctuations for facilitating electron transfer is given;
4. comparison of heterogeneous (electrochemical) and homogeneous rate constants;
5. distance effect on electron transfer rates; and
6. other aspects including orbital and orientation effect on electron transfer rates (preliminary calculations), photoelectric emission experiments, and application of the quadratic free energy expression to the other classes of reactions.

1. Electron Transfer Matrix Elements

Second order bimolecular rate constants k_{bi} have been expressed in terms of first order constants $k(R)$ at a reactants' separation distance R by integrating over all R using the pair distribution function $g(R)$, e. g., refs. (1-4):

$$k_{bi} = \int_0^{\infty} k(R)g(R) 4\pi R^2 dR \quad (1)$$

$g(R)$ can be expressed in terms of some "work" term $w_r(R)$ to bring

the reactants r to a separation distance R :

$$g(r) = \exp[-w_r(R)/kT] \quad (2)$$

Equation (1) is an approximation which tacitly assumes (4, 5) that R itself is not the reaction coordinate. With this approximation eq. (1) can be deduced from Ref. (5).

A first order nonadiabatic electron transfer rate constant $k_{na}(R)$ has been written in the form (1, 6)

$$k_{na}(R) = \frac{2\pi}{\hbar} |H_{12}|^2 \sum_{i,f} P_{f-i} p_i^{eq} \delta(E_f - E_i) \quad (3)$$

where P_{f-i} is the Franck-Condon overlap factor $|\langle \psi_f | \psi_i \rangle|^2$ of the wavefunctions for nuclear motion of reactants and solvent (in state i) and of products and solvents (in state f), p_i^{eq} is the Boltzmann distribution function for finding the system in the i 'th initial state of the entire system containing the reactants, and the Dirac delta function $\delta(E_f - E_i)$ ensures that there is conservation of energy during the electron transfer act. Because of the 'continuum' of final states f , the summation over f becomes an integral and the delta function disappears on integration.

We note that the maximum value that the double sum in eq. (3) can have is when there is extensive overlap of each ψ_i with a corresponding ψ_f (hence, no Franck-Condon barrier), so that for every i P_{f-i} is unity for some f . The double sum is then found to be (Appendix A), approximately $1/\hbar\omega$, where $\omega/2\pi$ is a typical relevant vibration frequency. Thus,

$$k_{na}^{max}(R) \approx \frac{2\pi}{\hbar} |H_{12}|^2 \frac{1}{\hbar\omega} \quad (4)$$

If k_{na}^{max} exceeds 10^{13} s^{-1} the nonadiabatic approximation (and hence eq. 4) breaks down, and the reaction is adiabatic under such conditions of no Franck-Condon barrier.

Some recent calculations of H_{12} for the $\text{Fe}(\text{H}_2\text{O})_6^{2+}$ reaction have been reported by Newton and co-workers (3, 7, 8). Previously, in calculations on this reaction a value of H_{12} of 2.7 cm^{-1} at $R = 7.3 \text{ \AA}$ had been obtained, but a later calculation showed that for a more elaborate wavefunction H_{12} at 7.3 \AA was 9 cm^{-1} , while with a more favorable geometry for overlap it became 26 cm^{-1} (8). The r_{max} which maximized the integrand in eq. (1) was roughly 5.3 \AA , and here the full calculation gave $H_{12} \sim 120$ to 130 cm^{-1} (8). An H_{12} of only $\sim 60 \text{ cm}^{-1}$ would have been sufficient to make k_{na}^{max} reach the adiabatic value ($\sim 10^{13} \text{ s}^{-1}$) had there been no Franck-Condon barrier. However,

the actual reaction is not quite adiabatic: adiabaticity also depends on the slopes of the reactants' and products' potential energy curves for the prevailing Franck-Condon barrier (7). (Cf. Landau-Zener theory for curve crossing problems.) The net result is that the reaction deviates from adiabaticity, in these available calculations, by a factor of about 1/5 (7). An important feature of attaining a substantial H_{12} in such calculations was the mutual penetration of the two aquo ions (i.e., $r \approx 5.3 \text{ \AA}$) or a suitable orientation. (However, the H_{12} 's may be inaccurate, because of the sensitivity of the tails of the wavefunctions to approximations.)

For reactants such as ruthenium bipyridyls or iron phenanthrolines, even some delocalization of the electron of the metal cation over the aromatic rings is expected to lead to a substantial H_{12} . Indeed, reactions of these compounds with other reactants, when accompanied by large negative ΔG° 's are diffusion-controlled, and so can't be far from adiabatic: If they were highly nonadiabatic they could never be diffusion-controlled.

2. Absolute Rate Calculations

The factors which influence the reaction rate constant include (i) the changes, as a result of electron transfer, within the innermost coordination shell of each reactant, (ii) the changes outside of this shell, (iii) the standard free energy of reaction (which is zero for self-exchange reactions), (iv) the frequency of "collisions", and (v) the electronic matrix element, when the reaction is predominantly nonadiabatic. The 'collision frequency' includes, in the electron transfer case, the range of R 's over which electron transfer can occur significantly (more precisely which contribute to the integral in eq. (1) (4)). We comment in Appendix B on the first factor.

Recently, there have been determinations of the metal-ligand bond lengths of electron transfer reactants in solution using EXAFS. This technique not only extends the X-ray data from the solid state to solutions but also increases considerably the number of systems for which the bond length changes Δq^0 are known. An extensive table is given by Brunschwig et al. (9). For a number of systems Δq^0 is very small: $\text{Fe}(\text{CN})_6^{3-,4-}$ (0.01 \AA), $\text{Fe}(\text{phen})_3^{2+,3+}$ (0.00), $\text{Co}(\text{bpy})_3^{2+,3+}$ (-0.02), $\text{Ru}(\text{NH}_3)_6^{2+,3+}$ (0.03). For such systems, the contribution of the innermost coordination shell to the reorganization energy is very small indeed, and is dominated by the solvent contribution from outside that shell. For a number of other systems, there is a substantial Δq^0 , and then the innermost coordination shell is a major contributor to the reaction barrier: $\text{Fe}(\text{H}_2\text{O})_6^{2+,3+}$ (0.13), $\text{Co}(\text{NH}_3)_6^{2+,3+}$ (0.22), $\text{Co}(\text{phen})_3^{2+,3+}$ (0.19), and $\text{Cr}(\text{H}_2\text{O})_6^{2+,3+}$ (0.32 for five ligands and 0.09 for one) (9). Nuclear tunneling factors usually contribute only between a factor of 1 to 7 in the self-exchange rate constants (10, 11). The results for the absolute rates, obtained with expressions which are substantially the same as those which we gave earlier (5, 12), yielded reasonable agreement with the experimental rate constants (13), considering that an error in the calculated free energy of

activation of $1.4 \text{ kcal mol}^{-1}$ yields an error of a factor of ten in the rate constant. Adiabaticity was assumed.

Although we have considered the rate constants themselves in this section, the most widely used application of the electron transfer theory we've described continues to be the 'cross-relation' (14),


$$k_{12} \sim (k_{11}k_{22}k_{12}f_{12})^{\frac{1}{2}} \quad (5)$$

relating the rate constants of cross-reactions to those of self-exchange reactions. A sampling of recent tests is given elsewhere (15). Tests involving cross reactions of inorganic reagents with cytochrome c was given by Sutin (16).

3. Solvent Effects

Several systems have been studied in which the $1/D_{op} - 1/D_s$ solvent effect (17) applies to electron transfer rate constants. These include the work on aromatic anions-aromatic molecules (18), bis-arene chromium $^{1+},^0$ reaction (19), the tris-hexafluoroacetylacetonato $Ru^{1+},^{2+}$ one (20), and various intervalence-transfer optical systems, e. g., (21-23). On the other hand, there was no correlation in the case of the ferrocene-ferrocenium ion $^0,^{1+}$ reaction (24) or in the case of the spectrum of compound II below (23).

The solvent effect represented by $1/D_{op} - 1/D_s$ is nonspecific and would be expected to break down when (i) the electronic overlap is no longer weak, or (ii) the solvent enters the the inner coordination shell, if any, or (iii) there is a specific solvent effect. An example of (i) is found in the pyrazine bridge system (compound II below). Results on charge transfer spectra have been reported (23) for I and II:



The results for I showed the expected (25) theoretical dependence of spectral absorption maximum on $(1/D_{op} - 1/D_s)$, with a slope which agreed with the theoretical value (D_{op} and D_s are the optical and static dielectric constants). Compound II showed no dependence--reflecting a strong rather than weak electronic interaction of the two Ru's via the pyrazine bridge.

Kosower and Huppert recently studied solvent effects on fluorescence quenching when the latter was caused by intramolecular charge transfer (26). They considered an amino naphthalenesulphonic acid derivative and a cyanodimethylamino benzene. In each case they found that the fluorescent lifetimes in a variety of solvents equaled the 'constant charge' longest dielectric relaxation time $\tau' = (D_{op}/D_s)\tau$ for each solvent, where τ is the usual ('constant field') relaxation time. For water, when $\tau \sim 8$ ps, $\tau' \sim 0.2$ ps. These results are striking in showing quite directly how experimental reaction rate constants can be related to bulk dielectric properties, in the present case when the reaction is 'barrierless'.

One can use this result to calculate the rate constant for formation of polarization fluctuations in a solvent: Suppose that τ' is the relaxation time in the sense of being the reciprocal of a rate constant (27) rather than being a 'half-life'. Let the fluctuation be such that its free energy of formation is ΔG^* . The relaxation time for decay of the fluctuation is $(D_{op}/D_s)\tau$, corresponding to a first order rate constant of $(D_s/D_{op}\tau)$. Thus, by microscopic reversibility the rate constant for forming the solvent fluctuation is

$$k = (D_s/D_{op}\tau)e^{-\Delta G^*/RT} \quad (6)$$

The value of ΔG^* for fluctuations in the solvent outside the innermost coordination shell has been given previously (17) in terms of $1/D_{op} - 1/D_s$. τ' has also been used in reaction rates for other types of reactions, e. g., (28).

4. Electrochemical Rate Constants

A recent survey of the comparison between homogeneous self-exchange rate constants k_{ex} and electrochemical rate constants k_{el} has been given by several authors (29, 30). The slope of the $\log k_{el}$ vs $\log k_{ex}$ is approximately $\frac{1}{2}$, as predicted for the case where the reactant center-reactant center distance in the homogeneous case is twice the reactant center-electrode surface distance (14). The very fast reactions showed enhanced deviations. The electrochemical rate constants in general tended to be somewhat slower than expected from the simple theory (roughly a factor of 10 in the region where the log-log plot had a slope of $\frac{1}{2}$). The reason may lie in a somewhat larger reactant-metal distance, resulting in a somewhat larger solvent reorganization energy and, possibly, in an extra nonadiabaticity.

The slope of the $\log k_{el}$ vs the activation overpotential plot has been determined for a number of organic reactants by Savéant and co-workers (31). The results show a curvature, as expected from theory.

5. Distance Effects in Electron Transfer Rates

At large separation distances the overlap of the electronic orbitals of the reactants becomes very small and the electronic H_{12} matrix element thereby also becomes small, varying roughly exponentially with separation distance R , and the reaction is now non-adiabatic. We have

$$k \propto e^{-\alpha R} \quad (7)$$

In frozen media the closer reactants react first and then the more distant ones. There results a peculiar kinetics for such second order reactions, the amount reacted now varying with $\log t$ (32-34). From such studies values of α have been estimated. For reactions between aromatic molecules and aromatic anions it is about 1.1 \AA (33), and between pyrene⁺ and tetramethylene phenylene diamine it is about 1.15 \AA^{-1} (34). Intramolecular electron transfer studies are desirable and are in progress in several laboratories on the effect of separation distance R on the rate.

We consider next a biological system, cytochrome *c*-cytochrome *c* peroxidase. A recent approximate estimate of the edge-to-edge distance between the hemes is about 16.5 \AA (35). The rate constant for electron transfer is known to be at least as fast as 10^4 s^{-1} (36). The maximum k expected if α were 1.1 \AA^{-1} would be $10^{13} \exp(-18.1)$, i. e., $\sim 10^5 \text{ s}^{-1}$, which, thus far, is consistent with the above existing minimum value of k .

A solvational distance effect, expected from the solvent reorganization theory (17), has been used to interpret some experimental results (37).

6. Other Aspects

Recent interesting photoelectric emission experiments and interpretation have been performed by Delahay and his group (38). He has been able to interpret the results in terms of concepts closely related to those used for thermal electron transfers and described earlier: contributions from the inner coordination shell and from the solvent outside it.

One continuing interest has been the investigation of highly exothermic reactions and a possible 'inverted' effect on the rate. We have discussed recent results elsewhere (39). A detailed calculation was presented (40) for the highly exothermic reaction of electronically-excited $\text{Ru}(\text{bipy})_3^{2+}$ with various metal bipyridyls. The absence of any large inverted effect in these reactions was clarified in part. However, not included was the role, if any, of high frequency CH vibrations in accepting some of the excess energy of reaction by producing excited CH vibrational states, and thereby reducing the

effective $-\Delta G^\circ$ acting on the other coordinates. Indeed, it would be useful to repeat these and related experiments using deuterated systems, which would have less ability to accept the excess energy of reaction and so might show a larger tendency to 'inverted' behavior if CH vibrations play a role.

Orientational or geometrical effects on electron transfer rates are known. It may be recalled that in the electron transfer reaction between hexahelicene and its anion, the df pair reacted four times more slowly than the dd pair, which itself was diffusion controlled (41). Orientational effects for spheroidal shaped molecules have been examined theoretically by Siders and Cave in our group using a simple model (constant potential energy inside the spheroid) which will be described elsewhere. Appropriate orientations produce a more favorable H_{12} . Larsson has discussed from a molecular orbital viewpoint orbital effects on electron transfer rates (42).

Reviews or adaptations of electron transfer theory to organic systems has been described by Ebersson (43) and by Kochi (44). The quadratic relation between the free energy of activation and that of reaction, first obtained for electron transfer reactions (17), has now been applied to a variety of nonelectron transfer organic systems, most recently by Murdoch (45).

III. Atom and Proton Transfers

We describe here some recent developments in light atom transfers for collinear collisions, and then extend the basic idea to actual systems.

In the transfer of an H between two heavy atoms, e. g.,



one expects that the H atom coordinate can be treated as 'fast' and the motion of the I's as 'slow'. (I. e., one has a Born-Oppenheimer type of approximation.) In the theory one would solve for the vibrational eigenvalue of the light particle (H), at fixed positions of the heavy particles I and then allow the I's to move in the presence of this effective potential created by the averaged motion of the H. To treat this in a practical way, Babamov and I used polar coordinates (R, θ) in the usual mass-weighted skewed-axes coordinate space (46). The radial coordinate R corresponds approximately to the I-I motion and the angular coordinate θ to the H motion.

Reaction (8) is a symmetric reaction, and was treated by Babamov in terms of symmetric (s) and antisymmetric (a) wavefunctions for the vibrational motion of the H in the presence of the I's (46). Since the H is initially localized on one of the I's, one can

expect that the probability of reaction will depend both on the s- and a- solutions. E. g., when R is large, an H-wavefunction localized near one I is approximately $\psi_s + \psi_a$, and when localized near the other it is approximately $\psi_s - \psi_a$.

The approach of the two I's under the influence of the symmetric state H-motion can be treated quantum mechanically, and similarly for the approach when the H-motion is in an antisymmetric vibrational state. One finds that the reaction probability p for a system in an initial HI vibrational state i to yield a final vibrational state i of the product IH is, approximately, (46)

$$p_{i \rightarrow i} = \sin^2(\eta_a - \eta_s) \quad (9)$$

where η_a (η_s) is the so-called phase shift for the I-I motion when the I's approach under an effective a- (s-) state of the H. By calculating these phase shifts, p was calculated and compared with the exact (many coupled states) quantum mechanical value for p, for the assumed potential energy surface. The results are given in Fig. 1, taken from a paper with Babamov and Lopez (47). The circles denote the exact quantum values (48) and the solid line is that calculated




Fig. 1 Reaction probability for reaction (8) when the reactants are in their lowest vibrational state, plotted versus total energy.

from eq (9), when the translational energy of relative motion of the I and HI is varied. Under the conditions investigated, p_{f-i} is negligible except for $f = i$, i. e., when the vibrational quantum number of the HI product equals that of the HI reactant.

The spike in Fig. 1 is due to a "resonance": the effective potential in the s- state of the H- motion forms a well for the I-I motion at smaller I-I distances, due to the delocalization of the H. This well creates a quasi-bound level, i. e., a "resonance" at some energy of the I-I motion and creates the spike. Equation (9) also has an approximate version (46) where

$$\eta_a - \eta_s \cong \gamma e^{-Q_0} \quad , \quad (10)$$

Q_0 being the usual tunneling integral for the H-motion at the distance R_0 of closest approach and γ a known constant. In this approximate form eq. (9) was applied by Bondi *et al.* to calculate rate constants for collinear collisions, with quite good agreement with exact numerical quantum values (49, 50).

The comparison in Fig. 1 shows that the behavior of the H atom transfer for this system is well understood, both qualitatively and quantitatively. For application to more complex systems, e. g., in solution, we note first that such systems will typically not be exactly "in resonance" as in eq. (9), when interactions with the environment are taken into account. Thus, for this reason a study was initiated of a nonsymmetric reaction



with $A \neq B$ (47). The reaction probability now depends on the energy defect ΔE between the effective potential energies of the (A, HB) and (AH, B) configurations (potential energy plus vibrational energy of the H motion) in the activated complex region ($\Delta E = 0$ in reaction (9)). In ref. (47a) an expression was given for the reaction probability which, in the threshold energy region (the region important for rate constants), can be written as (47b)

$$p_{f-i} = p_{f-i}^{\text{sym}} \exp[-\beta(\Delta E_{if})^2] \quad , \quad (12)$$

where p_{f-i}^{sym} is the same as for the symmetric reaction ($\Delta E = 0$).

ΔE_{if} is the ΔE for the ("diabatic", i. e., localized) (47) vibrational state i of (A, HB) and for state f of (AH, B), near the turning point of the R-motion, and β is a known constant. When β is sufficiently large the exponential behaves like the delta function that occurs in electron transfer reactions. ΔE_{if} was assumed small relative to vibrational spacings of E_f .

To apply (12) to an H-transfer in complex systems, we proceed as follows: We classify all coordinates as either fast or slow, the

latter including reaction coordinate. The 'fast' coordinates are taken to be the H being transferred and any other nearby H's, for example any attached to CH₃'s, NH₂'s, etc. which merely change slightly their equilibrium position or vibration frequencies, as well as any which may simultaneously transfer from one molecule to an adjacent one. Holding all remaining nuclei fixed, the energy defect is now written as $\Delta E_{if}(q)$, where i now denotes the (diabatic) vibrational state of all fast coordinates in the reactants. f is that in the products, and q denotes the totality of all slow coordinates. One of these q 's is designated as the reaction coordinate q_R , e. g., the AB distance in reaction (11), and the remaining q 's are denoted by q' . The reaction probability depends on the energy defect in the transition state region $\Delta E_{if}(q')$ (namely, at the turning point of the R-motion); the reaction probability also depends on the initial energy along q_R , $E_R(q_R)$, and on the initial state i of the H's. If we denote this reaction probability by p_{f-i} , the rate constant k is given by (51)

$$k = \sum_i \int \dots \int p_{f-i} p_{iq'}^{eq} dq' dE_R/h \quad (13)$$

where $p_{iq'}^{eq}(E_R) dq' dq_R dp_R/h$ is the equilibrium probability at the given temperature of finding the slow coordinates in the volume element dq' , the fast coordinates in a collective quantum state i , and the system in a phase space volume element $dq_R dp_R$ for the reaction coordinate. The integration is over all q' and E_R , and the sum is over all initial states i .

The detailed theory thus includes a calculation of the effective potential energies for the q_R motion, as a function of q , and of using a suitable expression for the reaction probability p . Should any of the q' 's in eq. (13) be, instead, high frequency motions, a quantum treatment for their motion can be used.

When there are additional H's as in nearby CH₃'s, NH₂'s, etc., expression for p_{f-i}^{sym} in (47) is modified; Franck-Condon factors for these additional H's could be added as a first approximation. (When there is a concerted transfer of H's, still further modification of (47) is needed, of course.)

Purely to illustrate eqs. (12) and (13) we give a calculation for a very simple model in which the effect of the q' 's on E_i and E_f is quadratic:

$$E_i(q') = E_i^0 + \frac{k_r}{2} (q' - q'_{ro})^2 \quad (14)$$

$$E_f(q') = E_f^0 + \frac{k_p}{2} (q' - q'_{po})^2 \quad (15)$$

(For notational brevity we have only used one term; k is actually a square matrix, one $q' - q'_{ro}$ is a column vector and the other a row vector.) In eq. (14) E_i is the value of E_i at some value of q' , $q' = q'_{ro}$, etc. If the additional contribution of the q' coordinates to the potential energy is written as $V_r(q')$ in the reactants' configuration, the integral over q' becomes

$$\int e^{-\beta(\Delta E_{if})^2} p_{iq}^{eq} dq' = \frac{\int e^{-\beta[\Delta E_{if}(q')]^2} e^{-V_r(q')/kT} e^{-E_i(q')/kT} dq'}{\sum_i \int e^{-V_r(q')/kT} e^{-E_i(q')/kT} dq'} \quad (16)$$

where $E_i(q')$ is given as a function of q' by eq. (14). For various models of $V_r(q')$ eq. (16) can be integrated. The presence of the $\beta(\Delta E_{if})^2$ term produces the type of reorganization which appeared in electron transfer reactions (5, 12, 17), when ΔE_{if} depends on q' . If it were independent of q' , the right hand side of eq. (16) would become simply $\exp[-\beta(\Delta E_{if})^2]$ and the result would reduce to that in ref. (47).

We plan to implement eq. (13) elsewhere for H and H^+ transfer. We discuss its relation to standard expressions in Sec. IV.

Finally, we comment briefly on a Franck-Condon type calculation for atom transfers. The profile of the potential energy surface vs the coordinate of the H-atom in transit is typically a double well. The latter has sometimes been approximated by two harmonic oscillator potentials, each localized on one of the two wells. Franck-Condon type arguments, e. g., use of eq. (3) involving the overlap of harmonic oscillator vibrational wavefunctions, are then used. However, such a procedure leads to orders of magnitude error in the absolute rates (52, 53). Thus, a different approach is desirable for atom transfers, such as that given in eq. (9) of the present paper for resonant symmetric on-resonance reactions and eq. (12) for asymmetric or off-resonant ones.

IV. Similarities and Differences Between Electron and H-Transfers

The principal similarity of these two types of reactions is the presence of fast and slow coordinates. In an electron transfer reaction the motion of the electrons in the reactants, and that in the surrounding solvent (appearing in the D_{op}) are the fast coordinates. The nuclei are the slow coordinates. In an H atom or H^+ transfer, the fast coordinates are those of all the relevant H's, including the H or H^+ 's being transferred and all remaining nearby H or H's. The slow coordinates are the rest of the coordinates (the q 's in the previous section).

This theory involving the assumption of both fast and slow coordinates may be compared or contrasted with the conventional approach in transition state theory. Here, it is often (though not always) assumed that all coordinates but the reaction coordinate are the fast coordinates. This approximation is made tacitly when one assumes that at each q_R the averaged motion of the other coordinates yields eigenvalues $E_i(q_R)$, which act as an effective potential for the motion along q_R . Such an approximation implies that all such coordinates are 'fast', relative to the reaction coordinate. One then focuses attention on these energy levels in the transition state region (at $q_R = q_R^\ddagger$) and calculates a rate constant. For H-transfers the tunneling integral used is the same as the WKB tunneling integral in Q_0 in eq. (10). What is new is eq. (9), the treatment using coupled H's (nearby H's), the presence of the γ (probably of the order of unity) in eq. (10), and the presence of the $\exp[-\beta(\Delta E_{if})^2]$ in eq. (12). The latter leads, as already noted, to a type of reorganizational term of the environment, neglected in conventional transition state theory. Another treatment of a reorganizational term, based on eq. (3) plus added approximations, rather than on eqs. (12)-(13), has been given by Dogonadze, Kuznetsov and Levich (54).

Equations (9), (10) and (12), when applied via eq. (13), appears to provide a dynamical treatment more detailed and fundamental than existing treatments of H and H^+ transfers. Having said this, one cannot overemphasize the detailed insight which conventional transition state theory has yielded for H and H^+ -transfers including, in the electrochemical case, the hydrogen evolution reaction. Equations (9), (10), (12) and (13) are intended to be applied to such reactions.

While electron and H-atom (proton) transfers have much in common, as we have noted above, there are also a number of differences:

(i) The Born-Oppenheimer type separation of variables is more accurate for electron-nuclei systems than for H atom-heavy nuclei systems. Thereby, the reaction probability is not as sharply a peaked function of the energy-defect ΔE in the transition state region. In electron transfer reactions the important role of this energy defect appears, in the case of the all-quantum treatment eq. (3), in the form of the Dirac delta function $\delta(E_f - E_i)$, while in a classical treatment it appears in the assignment of the transition state to the intersection of the reactants' and products' potential energy surfaces. In H transfers it appears in the present paper via an energy defect ΔE , as in eq. (12).

(ii) The H-atom transfer does not entail a charge transfer, and so does not involve the major solvent reorganization which accompanies electron transfer. The H^+ -transfer is frequently over a quite short distance (e. g., a transfer between adjacent molecules) and so it, too, is not expected under such conditions to yield a large solvent reorganization energy. Of course, when the proton transfer proceeds via an intervening water molecule $AH^+ \cdots OH_2 \cdots B$ and involves a

concerted motion of the two H^+ 's, the effective transfer of charge is over a larger distance, and so would entail a somewhat larger reorganization energy. Such a H_2O bridge removes any need, incidentally, for breaking any pre-existing H-bonds. However, in H^+ transfers involving carbon acids and bases, unlike transfers solely between N's and O's, there may be some preliminary desolvation of the noncarbon acid or base, creating thereby an added barrier to the reaction in the form of a "work term" (55).

(iii) Electron transfers usually involve a weak overlap of the electronic orbitals of the reactions, whereas H atom transfers or H^+ transfers are expected to involve strong overlap. This difference requires a more detailed electronic structure calculation of the potential energy surface in the vicinity of the transition state for H- and H^+ -transfers than was necessary for electron transfers.

(iv) In electron transfers there are usually only two electronic states accessible--one for the reactants and one for the products--because of the wide spacing of electronic energy levels. (Reactions at metal electrodes are an exception, because of the many electronic levels in a metal (1, 5). No inverted effect is predicted to occur there, therefore (63). Again, in a homogeneous reaction an excited electronic state of the reaction products may also become accessible.) In the case of H's or H^+ 's, however, there are frequently a substantial number of vibrational states of the H's accessible, rather than just two.

(v) In sufficiently exothermic H or H^+ transfer reactions the reaction may be 'downhill' throughout, with no need for H-tunneling (56). A corresponding phenomenon is unlikely for electron transfers, except for 'strong overlap' reactions.

V. Summary

In the present paper we have summarized some recent experiments on electron transfer reactions, recent calculations based on the equations of ref. 5, apart from inclusion of a relatively minor nuclear tunneling factor or an 'electron transfer over a distance' factor. These two factors affect the rate in opposite directions. The calculation of the frequency of solvent fluctuations outside the innermost coordination shell of an ion was described in eq. (6).

We have also described a recent development in the dynamics of H-atom transfers due to Babamov, Lopez and Marcus. We have now outlined how to extend it to H or H^+ transfers in more complicated systems and in condensed media. The treatment is based on a Born-Oppenheimer type separation of fast coordinates (the nearby H's, including the one or two being transferred) and slow coordinates (all other coordinates). Similarities and differences with the theory

of electron transfer are compared in Sec. IV, and both are compared with conventional transition state theory.

Acknowledgment

It is a pleasure to acknowledge the support of this research by the Office of Naval Research. Contribution No. 6856.

Appendix A. Derivation of Equation (4)

To obtain k_{na}^{\max} in eq. (3) we first set the Franck-Condon factor P_{f-i} equal to its maximum value of unity. The sum over f is then replaced by an integral over E_f : $\int (df/dE_f) dE_f \delta(E_i - E_f)$, where df/dE_f is the density of final states. For a one-coordinate type problem (e. g., a concerted combination of outward symmetric breathing vibration of one reactant with inward breathing vibration of the other in an electron transfer), the density of final vibrational states is $1/\hbar\omega$ where $\hbar\omega$ is the vibrational quantum. Integration then yields eq. (4). Equation (3) assumes, of course, that the density of final states is sufficiently large that it can be treated as a continuum. Typically, the many degrees of freedom provide this continuum, but we have used a pure vibrational model for purposes of deriving eq. (4).

Appendix B. Inner Coordination Shell Contribution to the Energy Barrier

One interesting aspect of the calculations of the innermost coordination shell involves the use of appropriate force constants: If a detailed force constant analysis of the metal-ligand stretching vibrations is made, there are diagonal and off-diagonal bond force constants. Both terms contribute to the differences in the various normal mode metal-ligand stretching frequencies. In calculating the contribution of the innermost shell to the energy barrier, all of these constants are to be included. The calculation is made simpler, when the normal modes remain intact upon electron transfer, by using normal mode force constants (12). They are automatically diagonal but contain both the diagonal and off-diagonal bond force constants as contributing factors. Several recent calculations (57, 58) included only the diagonal bond force constants, as we and others have noted (59). This point of using the normal mode force constants was indeed stressed earlier (5) and used in an early numerical calculation (60). Other treatments used diagonal bond force constants (61). Hush (62) calculated potential energy surfaces for the innermost coordination shell, rather than using known vibration frequencies, and then used the equivalent of a small displacement approximation.

References

1. V. G. Levich, *Adv. Electrochem. Electrochem. Eng.* 4, 249 (1966); V. G. Levich and R. R. Dogonadze, *Coll. Czech. Chem. Commun.* 26, 193 (1961), Transl., O. Boshko, University of Ottawa, Ontario, Canada.
2. E. G., P. R. Butler, M. J. Pilling, S. A. Rice and T. J. Stone, *Can. J. Chem.* 55, 2124 (1977); J. Keizer, *J. Phys. Chem.* 85, 940 (1981).
3. M. D. Newton, *Int. J. Quant. Chem. Symp.* 14, 363 (1980).
4. R. A. Marcus, *Intl. J. Chem. Kinetics* 13, 865 (1981).
5. R. A. Marcus, *J. Chem. Phys.* 43, 679 (1965).
6. Cf. N. Kestner, J. Logan and J. Jortner, *J. Phys. Chem.* 78, 2148 (1974); see also D. Devault, *Quart. Rev. Biophys.* 13, 4 (1980), which contains a nice review of the various theories.
7. J. Logan and M. D. Newton, *J. Chem. Phys.* 78, 4086 (1983) and references cited therein.
8. H. L. Friedman and M. D. Newton, *Faraday Disc. Chem. Soc.* 74, 73 (1982).
9. B. S. Brunshawig, C. Creutz, D. H. Macartney, T.-K. Sham and N. Sutin, *Faraday Disc. Chem. Soc.* 74, 113 (1982).
10. B. S. Brunshawig, J. Logan, M. D. Newton and N. Sutin, *J. Am. Chem. Soc.* 102, 5798 (1980).
11. P. Siders and R. A. Marcus, *J. Am. Chem. Soc.* 103, 741 (1981); E. Buhks, M. Bixon, G. Navon and J. Jortner, *J. Phys. Chem.* 85, 3759 (1981).
12. R. A. Marcus, *Discuss. Faraday Soc.* 29, 21 (1960).
13. N. Sutin, *Progr. Inorg. Chem.* 30, 441 (1983).
14. R. A. Marcus, *J. Phys. Chem.* 67, 853, 2889 (1963).
15. R. A. Marcus and N. Sutin, *Biochim. Biophys. Acta* (to be submitted).
16. N. Sutin, *Adv. Chem. Ser.* 162, 156 (1977).
17. R. A. Marcus, *J. Chem. Phys.* 24, 966 (1956).
18. J. R. Brandon and L. M. Dorfman, *J. Chem. Phys.* 53, 3849 (1970).
19. T. T.-T. Li and C. H. Brubaker, Jr., *J. Organomet. Chem.* 216, 223 (1981).
20. M.-S. Chan and A. C. Wahl, *J. Phys. Chem.* 86, 126 (1982).
21. G. M. Tom, C. Creutz and H. Taube, *J. Am. Chem. Soc.* 96, 25 (1974).
22. R. W. Callahan and T. J. Meyer, *Chem. Phys. Lett.* 39, 82 (1976); R. W. Callahan, F. R. Keene, T. J. Meyer and D. J. Salmon, *J. Am. Chem. Soc.* 99, 1064 (1977); M. J. Powers and T. J. Meyer, *J. Am. Chem. Soc.* 100, 14, 4393 (1978); M. J. Powers and T. J. Meyer, *J. Am. Chem. Soc.* 102, 1289 (1980); B. P. Sullivan and T. J. Meyer, *Inorg. Chem.* 19, 752 (1980).
23. C. Creutz, *Progr. Inorg. Chem.* 30, 1 (1983); C. Creutz, *Inorg.*

- Chem. 17, 3723 (1978).
24. E. S. Yang, M. -S. Chan and A. C. Wahl, J. Phys. Chem. 84 3094 (1980).
25. N. S. Hush, Progr. Inorg. Chem. 8, 391 (1967); cf. R. A. Marcus, J. Chem. Phys. 43, 1261 (1965).
26. E. M. Kosower and D. Huppert, Chem. Phys. Lett. 96, 433 (1983).
27. H. Fröhlich, Theory of Dielectrics (Oxford, New York, 1949), p. 72.
28. G. van der Zwan and J. T. Hynes, J. Chem. Phys. 76, 2993 (1982); see also S. M. Mozumder, *ibid.*, 50 3153 (1969) and W. J. Chase and J. W. Hunt, J. Phys. Chem. 79, 2835 (1975).
29. T. Saji, T. Yamada and S. Aoyagui, J. Electroanal. Chem. 61, 147 (1975).
30. M. J. Weaver and J. T. Hupp, ACS Symp. Ser. 198, 181 (1982); M. J. Weaver, J. Phys. Chem. 84, 568 (1980).
31. J. M. Savéant and D. Tessier, Faraday Disc. Chem. Soc. 74, 57 (1982) and references cited therein.
32. M. Tachiya and A. Mozumder, Chem. Phys. Lett. 28, 87 (1974).
33. I. V. Alexandrov, R. F. Khairutdinov and K. I. Zamaraev, Chem. Phys. 32, 123 (1978).
34. J. R. Miller and J. V. Beitz, J. Chem. Phys. 74, 6746 (1981). α = their 2γ .
35. T. L. Poulos and J. Kraut, J. Biol. Chem. 255, 10322 (1980).
36. J. J. Leonard and T. Yonetani, Biochemistry 13, 1465 (1974).
37. M. J. Powers and T. J. Meyer, J. Am. Chem. Soc. 102, 1289 (1980); M. J. Powers, D. J. Salmon, R. W. Callahan and T. J. Meyer, *ibid.*, 98, 6731 (1976); A. P. Szecsy and A. Haim, J. Am. Chem. Soc. 103, 1679 (1981); A. Haim, Pure and Appl. Chem. 55, 89 (1983).
38. P. Delahay, Chem. Phys. Lett. 89, 149 (1982; *ibid.*, 90, 425 (1982); *ibid.*, (in press), and references cited therein.
39. R. A. Marcus, Faraday Disc. Chem. Soc. 74, 7 (1982).
40. R. A. Marcus and P. Siders, J. Am. Chem. Soc. 86, 622 (1982).
41. R. Chang and S. I. Weissman, J. Am. Chem. Soc. 89, 23 (1967).
42. S. Larsson, J. Am. Chem. Soc. 103, 4034 (1981); S. Larsson, Theoret. Chim. Acta (Berl.) 60, 111 (1981).
43. L. Ebersson, Adv. Phys. Org. Chem. 18, 79 (1982).
44. J. K. Kochi, Pure and Appl. Chem. 52, 571 (1980).
45. R. M. G. Roberts, D. Ostović and M. M. Kreevoy, Faraday Disc. Chem. Soc. 74, 257 (1982); W. J. Albery, Annu. Rev. Phys. Chem. 31, 227 (1980); J. R. Murdoch, Faraday Disc. Chem. Soc. 74, 297 (1982) and references cited therein (discussion comment). See also (52).
46. V. K. Babamov and R. A. Marcus, J. Chem. Phys. 74, 1790 (1981).
47. (a) V. K. Babamov, V. Lopez and R. A. Marcus, J. Chem. Phys. 78, 5621 (1983); (b) same authors, Chem. Phys. Lett. in press.
48. J. A. Kaye and A. Kuppermann, J. Chem. Phys. 77, 573 (1981).
49. D. K. Bondi, J. N. L. Connor, B. C. Garrett and D. G. Truhlar, J. Chem. Phys. 78, 5981 (1983).

50. Another application of the ideas in (46) and examination of the corresponding classical behavior is given in C. Hiller, J. Manz, W. H. Miller and J. Römelt, *J. Chem. Phys.* **78**, 3850 (1983).
51. A standard argument is used: the equilibrium probability $p_{iq}^{eq} dq' dq_R dp_R / h$ is divided by dq_R to obtain a linear density and multiplied by the velocity \dot{q}_R to obtain the incident flux; $\dot{q}_R dp_R$ is readily shown to be dE_R . Cf. R. A. Marcus, *J. Chem. Phys.* **45**, 2138 (1966), derivation of eq. (3) there; R. A. Marcus in *Techniques of Chemistry*, Vol. 6, Pt. 1, E. S. Lewis, ed., Chap. 2.
52. K. H. Fung and K. F. Freed, *Chem. Phys.* **30**, 249 (1978).
53. V. Lopez, V. K. Babamov and R. A. Marcus, (to be submitted).
54. R. R. Dogonadze, A. M. Kuznetsov, and V. G. Levich, *Electrochim. Acta*, **13**, 1025 (1968).
55. M. M. Kreevoy and D. E. Konasewich, *Adv. Chem. Phys.* **21**, 243 (1971); M. M. Kreevoy and S. W. Oh, *J. Am. Chem. Soc.* **95**, 4805 (1973); A. I. Hassid, M. M. Kreevoy and T.-M. Liang, *Faraday Symp. Chem. Soc.* **10**, 69 (1975).
56. R. A. Marcus, *Faraday Symp. Chem. Soc.* **10**, 60 (1975); R. A. Marcus, *Physicochemical Hydrodynamics: V. G. Levich Festschrift*, Vol. 1, ed. D. B. Spalding (Advance Publications, London, 1977), p. 473.
57. H. C. Stynes and J. A. Ibers, *Inorg. Chem.* **10**, 2304 (1971), a result used in E. Buhks, M. Bixon, J. Jortner and G. Navon, *ibid.*, **18**, 2014 (1979).
58. J. M. Hale in *Reactions of Molecules at Electrodes*, N. S. Hush, ed. (Wiley-Interscience, New York, 1971), p. 229 ff., a result
59. Regarding (57) see (11); regarding (58) see S. U. M. Khan and J. O'M. Bockris, *Chem. Phys. Lett.* **99**, 83 (1983). Cf. P. Delahay, *ibid.* **99**, 87 (1983).
60. R. A. Marcus, *Trans. N. Y. Acad. Sci.* **19**, 423 (1957).
61. P. George and J. S. Griffith, in *The Enzymes*, P. D. Boyer, H. Lardy and K. Myrback, eds. Vol. 1, (Academic Press, New York, 1959), p. 347; cf. L. E. Orgel, *Inst. Intern. Chim. Solvay, Conseil. Chim.*, Brussels **10**, 289 (1956), which gives a somewhat different result. References (60) and (61) limited their calculation of the innermost coordination shell to self-exchange reactions. In ref. (12) cross-reactions were also included.
62. N. S. Hush, *Trans. Faraday Soc.* **57**, 557 (1961). An explicitly harmonic version is given in *Electrochim. Acta* **13**, 1005 (1968) and in *Progr. Inorg. Chem.* **8**, 391 (1967).
63. R. A. Marcus, *J. Chem. Phys.* **43**, 2654 (1965).

Jan 84

DL/413/83/01
GEN/413-2TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 N. Quincy Street Arlington, Virginia 22217	2	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152	1
ONR Pasadena Detachment Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	Naval Weapons Center Attn: Dr. A. B. Amster Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1	Dean William Tolles Naval Postgraduate School Monterey, California 93940	1
Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Mr. Vincent Schaper DTNSRDC Code 2830 Annapolis, Maryland 21402	1
DTNSRDC Attn: Dr. G. Bosmajian Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1	Mr. A. M. Anzalone Administrative Librarian PLASTEC/ARRADCOM Bldg 3401 Dover, New Jersey 07801	1

TECHNICAL REPORT DISTRIBUTION LIST, 359

Dr. David Aikens
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. A. P. B. Lever
Chemistry Department
York University
Downsview, Ontario M3J1P3

Dr. Stanislaw Szpak
Naval Ocean Systems Center
Code 6343, Bayside
San Diego, California 95152

Dr. Gregory Farrington
Department of Materials Science
and Engineering
University of Pennsylvania
Philadelphia, Pennsylvania 19104

M. L. Robertson
Manager, Electrochemical
and Power Sources Division
Naval Weapons Support Center
Crane, Indiana 47522

Dr. T. Marks
Department of Chemistry
Northwestern University
Evanston, Illinois 60201

Dr. Micha Tomkiewicz
Department of Physics
Brooklyn College
Brooklyn, New York 11210

Dr. Lesser Blum
Department of Physics
University of Puerto Rico
Rio Piedras, Puerto Rico 00931

Dr. Joseph Gordon, II
IBM Corporation
K33/281
5600 Cottle Road
San Jose, California 95193

Dr. D. H. Whitmore
Department of Materials Science
Northwestern University
Evanston, Illinois 60201

Dr. Alan Bewick
Department of Chemistry
The University of Southampton
Southampton, SO9 5NH ENGLAND

Dr. E. Anderson
NAVSEA-56233 NC #4
2541 Jefferson Davis Highway
Arlington, Virginia 20362

Dr. Bruce Dunn
Department of Engineering &
Applied Science
University of California
Los Angeles, California 90024

Dr. Elton Cairns
Energy & Environment Division
Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

Dr. D. Cipris
Allied Corporation
P.O. Box 3000R
Morristown, New Jersey 07960

Dr. M. Philpott
IBM Corporation
5600 Cottle Road
San Jose, California 95193

Dr. Donald Sandstrom
Department of Physics
Washington State University
Pullman, Washington 99164

Dr. Carl Kannewurf
Department of Electrical Engineering
and Computer Science
Northwestern University
Evanston, Illinois 60201

TECHNICAL REPORT DISTRIBUTION LIST, 359

Dr. Paul Delahay
Department of Chemistry
New York University
New York, New York 10003

Dr. P. J. Hendra
Department of Chemistry
University of Southampton
Southampton SO9 5NH
United Kingdom

Dr. T. Katan
Lockheed Missiles and
Space Co., Inc.
P.O. Box 504
Sunnyvale, California 94088

Dr. D. N. Bennion
Department of Chemical Engineering
Brigham Young University
Provo, Utah 84602

Dr. R. A. Marcus
Department of Chemistry
California Institute of Technology
Pasadena, California 91125

Mr. Joseph McCartney
Code 7121
Naval Ocean Systems Center
San Diego, California 92152

Dr. J. J. Auburn
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Joseph Singer, Code 302-1
NASA-Lewis
21000 Brookpark Road
Cleveland, Ohio 44135

Dr. P. P. Schmidt
Department of Chemistry
Oakland University
Rochester, Michigan 48063

Dr. H. Richtol
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 44106

Dr. C. E. Mueller
The Electrochemistry Branch
Naval Surface Weapons Center
White Oak Laboratory
Silver Spring, Maryland 20910

Dr. Sam Perone
Chemistry & Materials
Science Department
Lawrence Livermore National Lab.
Livermore, California 94550

Dr. Royce W. Murray
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. G. Goodman
Johnson Controls
5757 North Green Bay Avenue
Milwaukee, Wisconsin 53201

Dr. B. Brummer
EIC Incorporated
111 Chapel Street
Newton, Massachusetts 02158

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Electrochimica Corporation
Attn: Technical Library
2485 Charleston Road
Mountain View, California 94040

Library
Duracell, Inc.
Burlington, Massachusetts 01803

Dr. A. B. Ellis
Chemistry Department
University of Wisconsin
Madison, Wisconsin 53706

TECHNICAL REPORT DISTRIBUTION LIST, 359

Dr. M. Wrighton
Chemistry Department
Massachusetts Institute
of Technology
Cambridge, Massachusetts 02139

Dr. B. Stanley Pons
Department of Chemistry
University of Utah
Salt Lake City, Utah 84112

Donald E. Mains
Naval Weapons Support Center
Electrochemical Power Sources Division
Crane, Indiana 47522

S. Ruby
DOE (STOR)
M.S. 6B025 Forrestal Bldg.
Washington, D.C. 20595

Dr. A. J. Bard
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. Janet Osteryoung
Department of Chemistry
State University of New York
Buffalo, New York 14214

Dr. Donald W. Ernst
Naval Surface Weapons Center
Code R-33
White Oak Laboratory
Silver Spring, Maryland 20910

Mr. James R. Moden
Naval Underwater Systems Center
Code 3632
Newport, Rhode Island 02840

Dr. Bernard Spielvogel
U.S. Army Research Office
P.O. Box 12211
Research Triangle Park, NC 27709

Dr. William Ayers
ECD Inc.
P.O. Box 5357
North Branch, New Jersey 08876

Dr. M. M. Nicholson
Electronics Research Center
Rockwell International
3370 Miraloma Avenue
Anaheim, California

Dr. Michael J. Weaver
Department of Chemistry
Purdue University
West Lafayette, Indiana 47907

Dr. R. David Rauh
EIC Corporation
111 Chapel Street
Newton, Massachusetts 02158

Dr. Aaron Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02192

Dr. Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton SO9 5NH ENGLAND

Dr. R. A. Osteryoung
Department of Chemistry
State University of New York
Buffalo, New York 14214

Dr. Denton Elliott
Air Force Office of Scientific
Research
Bolling AFB
Washington, D.C. 20332

Dr. R. Nowak
Naval Research Laboratory
Code 6130
Washington, D.C. 20375

Dr. D. F. Shriver
Department of Chemistry
Northwestern University
Evanston, Illinois 60201

Dr. Aaron Fletcher
Naval Weapons Center
Code 3852
China Lake, California 93555

TECHNICAL REPORT DISTRIBUTION LIST, 359

Dr. Robert Somoano
Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California 91103

Dr. Johann A. Joebstl
USA Mobility Equipment R&D Command
DRDME-EC
Fort Belvoir, Virginia 22060

Dr. Judith H. Ambrus
NASA Headquarters
M.S. RTS-6
Washington, D.C. 20546

Dr. Albert R. Landgrebe
U.S. Department of Energy
M.S. 68025 Forrestal Building
Washington, D.C. 20595

Dr. J. J. Brophy
Department of Physics
University of Utah
Salt Lake City, Utah 84112

Dr. Charles Martin
Department of Chemistry
Texas A&M University
College Station, Texas 77843

Dr. H. Tachikawa
Department of Chemistry
Jackson State University
Jackson, Mississippi 39217

Dr. Theodore Beck
Electrochemical Technology Corp.
3935 Leary Way N.W.
Seattle, Washington 98107

Dr. Farrell Lytle
Boeing Engineering and
Construction Engineers
P.O. Box 3707
Seattle, Washington 98124

Dr. Robert Gotscholl
U.S. Department of Energy
MS G-226
Washington, D.C. 20545

Dr. Edward Fletcher
Department of Mechanical Engineering
University of Minnesota
Minneapolis, Minnesota 55455

Dr. John Fontanella
Department of Physics
U.S. Naval Academy
Annapolis, Maryland 21402

Dr. Martha Greenblatt
Department of Chemistry
Rutgers University
New Brunswick, New Jersey 08903

Dr. John Wasson
Syntheco, Inc.
Rte 6 - Industrial Pike Road
Gastonia, North Carolina 28052

Dr. Walter Roth
Department of Physics
State University of New York
Albany, New York 12222

Dr. Anthony Sammells
Eltron Research Inc.
710 E. Ogden Avenue #108
Naperville, Illinois 60540

Dr. W. M. Risen
Department of Chemistry
Brown University
Providence, Rhode Island 02192

Dr. C. A. Angell
Department of Chemistry
Purdue University
West Lafayette, Indiana 47907

Dr. Thomas Davis
Polymer Science and Standards
Division
National Bureau of Standards
Washington, D.C. 20234

END

FILMED

3-84

DTIC